

BEHAVIOR OF MONOPOLE TRAJECTORIES IN THE INSTANTON AND ANTI-INSTANTON SYSTEM AT FINITE TEMPERATURE

F. Araki, H. Suganuma and H. Toki
Research Center for Nuclear Physics, Osaka University
Mihogaoka 10-1, Ibaraki, Osaka 567, Japan

We study the connection between monopoles and instantons in the Polyakov-like gauge, and the behavior of the monopole trajectories generated by instantons at finite temperature. The monopole trajectories become long and complicated at the zero-temperature which can be regarded as monopole condensation. On the other hand, they tend to become simple at finite temperature. It is suggested that such a change is related to the confinement-deconfinement phase transition.

1 Introduction

The nonabelian gauge theory like QCD is reduced to the abelian gauge theory including monopoles under the abelian gauge fixing¹. It is thought that monopoles play a relevant role of the nonperturbative phenomena of QCD like the color confinement through monopole condensation. In the abelian gauge, the monopole appears from the hedgehog-like gauge configuration^{1,2}. On the other hand, the nonabelian gauge theory in the Euclidean space has a classical solution called instanton. Instantons are also believed to play an important role in the QCD ground state.

In the abelian-dominant system, instantons seem to lose the topological basis for its existence, since the instanton is a topological object in the non-abelian manifold. However, even in the abelian gauge, nonabelian components are not completely suppressed around the topological defect, *i.e.* monopoles. Therefore instantons are expected to appear only around the monopole trajectories in the abelian-dominant system. In fact, the fourth component of the exact instanton solution in the singular gauge becomes the hedgehog configuration as

$$A_4 = \frac{i}{\phi} \sum_k \frac{a_k^2 \sigma \cdot (\mathbf{x} - \mathbf{x}_k)}{|\mathbf{x} - \mathbf{x}_k|^4} \quad , \quad \phi \equiv 1 + \sum_k \frac{a_k^2}{|\mathbf{x} - \mathbf{x}_k|^2}, \quad (1)$$

where x_k is the k -th instanton center position and a_k is its size. The Polyakov-like gauge^{3,4,5,6}, in which A_4 is diagonalized, provides a monopole on $\mathbf{x} = \mathbf{x}_k$. Thus, the instanton center is inevitably penetrated by the monopole trajectory along the temporal direction in the Polyakov-like gauge.

2 Instanton and anti-instanton system at finite temperature

The exact solution of the Yang-Mills theory regards either instantons or anti-instantons. However, since it seems natural that QCD has both of them, we choose the R-ansatz⁷,

$$A^\mu = -i\sigma^a \cdot \frac{\eta_-^{a\mu\nu}\partial_\nu\phi_I + \eta_+^{a\mu\nu}\partial_\nu\phi_A}{\phi_I + \phi_A - 1} \quad (2)$$

with $\eta_\pm^{a\mu\nu} \equiv \varepsilon^{a\mu\nu 4} \pm \delta^{a\mu}\delta^{\nu 4} \mp \delta^{a\nu}\delta^{\mu 4}$. Here, ϕ_I and ϕ_A give instanton and anti-instanton profiles, respectively. These are the same form as ϕ of Eq. (1). The finite temperature system is described by imposing the periodic boundary condition on the Euclidean temporal direction, with the period $2\pi T$, where T is the temperature of the system. The profile function ϕ is modified as

$$\phi \longrightarrow 1 + \pi T \sum_k \frac{a_k^2}{r_k} \cdot \frac{\sinh(2\pi T r_k)}{\cosh(2\pi T r_k) - \cos(2\pi T \tau_k)}, \quad (3)$$

with $r_k \equiv |\mathbf{x} - \mathbf{x}_k|$ and $\tau_k \equiv t - t_k$. In the Polyakov-like gauge, the monopole trajectories appear on each center of the instanton and anti-instanton as the case of the previous section.

3 Numerical Simulation

Next, we demonstrate the behavior of the monopole trajectories in the above system by numerical simulations. In Figs. 1 and 2, we show examples of the monopole trajectories in this system at the temperature $T = 0$ and $T = 1(\text{fm}^{-1})$, respectively. In both cases, the monopole trajectories are generated by 100 instantons and 100 anti-instantons which are put on the zt -plane. The black dot and the small circle denote the instanton and anti-instanton center, respectively. In Fig. 1, the monopole trajectories are found to be highly complicated penetrating the centers of the instantons and the anti-instantons. As the temperature goes high (Fig. 2), the monopole trajectories tend to become straight lines along the temporal direction and change their topology.

4 Summary

We have studied the connection between monopoles and instantons. In the Polyakov-like gauge, monopoles appear on each instanton or anti-instanton center. Therefore, the monopole trajectories inevitably penetrate through instanton centers. We have studied the behavior of the monopole trajectories in

the instanton and anti-instanton system at finite temperature. At the zero-temperature, the monopole trajectories tend to become highly complicated and very long, which can be regarded as a signal of monopole condensation. When the temperature goes high, the monopole trajectories tend to become simple, and change their topology, which may correspond to the vanishing of monopole condensation.

Figure 1: The monopole trajectories in the instanton and anti-instanton system at the temperature $T = 0$. The monopole trajectories are found to be highly complicated penetrating through the centers of the instantons and the anti-instantons.

Figure 2: The monopole trajectories at $T = 1(\text{fm}^{-1})$. As the temperature goes high, the monopole trajectories tend to become straight lines along the temporal direction.

References

1. G. 't Hooft, *Nucl. Phys. B* **190**, 455 (1981).
2. H. Suganuma, S. Sasaki and H. Toki, *Nucl. Phys. B* **435**, 207 (1995).
3. H. Suganuma, K. Itakura, H. Toki and O. Miyamura, in *Nonperturbative Approaches to QCD*, 100 (PNPI, 1995).
4. H. Suganuma, S. Sasaki, H. Ichie, H. Toki and F. Araki, in *Frontier 96*, 177 (World Scientific, 1996).
5. F. Araki, H. Suganuma and H. Toki, in *RCNP Annual Report 1995*, 81 (1996).
6. H. Suganuma, S. Sasaki, H. Ichie, F. Araki and O. Miyamura, *Nucl. Phys.(Proc. Suppl.) B* **53**, 528 (1997).
7. E. V. Shuryak, *Nucl. Phys. B* **302**, 574 (1988).



